

20. COSMIC RAYS

Revised February 2000 by T.K. Gaisser and T. Stanev (Bartol Research Inst., Univ. of Delaware).

20.1. Primary spectra

The cosmic radiation incident at the top of the terrestrial atmosphere includes all stable charged particles and nuclei with lifetimes of order 10^6 years or longer. Technically, “primary” cosmic rays are those particles accelerated at astrophysical sources and “secondaries” are those particles produced in interaction of the primaries with interstellar gas. Thus electrons, protons and helium, as well as carbon, oxygen, iron, and other nuclei synthesized in stars, are primaries. Nuclei such as lithium, beryllium, and boron (which are not abundant end-products of stellar nucleosynthesis) are secondaries. Antiprotons and positrons are partly, if not entirely, secondaries, but the fraction of these particles that may be primary is a question of current interest.

Apart from particles associated with solar flares, the cosmic radiation comes from outside the solar system. The incoming charged particles are “modulated” by the solar wind, the expanding magnetized plasma generated by the Sun, which decelerates and partially excludes the lower energy galactic cosmic rays from the inner solar system. There is a significant anticorrelation between solar activity (which has an eleven-year cycle) and the intensity of the cosmic rays with energies below about 10 GeV. In addition, the lower-energy cosmic rays are affected by the geomagnetic field, which they must penetrate to reach the top of the atmosphere. Thus the intensity of any component of the cosmic radiation in the GeV range depends both on the location and time.

There are four different ways to describe the spectra of the components of the cosmic radiation: (1) By particles per unit rigidity. Propagation (and probably also acceleration) through cosmic magnetic fields depends on gyroradius or *magnetic rigidity*, R , which is gyroradius multiplied by the magnetic field strength:

$$R = \frac{pc}{Ze} = r_L B . \quad (20.1)$$

(2) By particles per energy-per-nucleon. Fragmentation of nuclei propagating through the interstellar gas depends on energy per nucleon, since that quantity is approximately conserved when a nucleus breaks up on interaction with the gas. (3) By nucleons per energy-per-nucleon. Production of secondary cosmic rays in the atmosphere depends on the intensity of nucleons per energy-per-nucleon, approximately independently of whether the incident nucleons are free protons or bound in nuclei. (4) By particles per energy-per-nucleus. Air shower experiments that use the atmosphere as a calorimeter generally measure a quantity that is related to total energy per particle.

The units of differential intensity I are $[\text{cm}^{-2}\text{s}^{-1}\text{sr}^{-1}\mathcal{E}^{-1}]$, where \mathcal{E} represents the units of one of the four variables listed above.

The intensity of primary nucleons in the energy range from several GeV to somewhat beyond 100 TeV is given approximately by

$$I_N(E) \approx 1.8 E^{-\alpha} \frac{\text{nucleons}}{\text{cm}^2 \text{ s sr GeV}} , \quad (20.2)$$

2 20. Cosmic rays

where E is the energy-per-nucleon (including rest mass energy) and $\alpha (\equiv \gamma + 1) = 2.7$ is the differential spectral index of the cosmic ray flux and γ is the integral spectral index. About 79% of the primary nucleons are free protons and about 70% of the rest are nucleons bound in helium nuclei. The fractions of the primary nuclei are nearly constant over this energy range (possibly with small but interesting variations). Fractions of both primary and secondary incident nuclei are listed in Table 20.1. Figure 20.1 [1] shows the major components as a function of energy at a particular epoch of the solar cycle. There has been a series of more precise measurements of the primary spectrum of protons and helium in the past decade [2,3,4,5].

Table 20.1: Relative abundances F of cosmic-ray nuclei at 10.6 GeV/nucleon normalized to oxygen ($\equiv 1$) [6]. The oxygen flux at kinetic energy of 10.6 GeV/nucleon is $3.26 \times 10^{-6} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1} (\text{GeV/nucleon})^{-1}$. Abundances of hydrogen and helium are from Ref. 2.

Z	Element	F	Z	Element	F
1	H	485	13–14	Al-Si	0.19
2	He	26	15–16	P-S	0.03
3–5	Li-B	0.40	17–18	Cl-Ar	0.01
6–8	C-O	2.20	19–20	K-Ca	0.02
9–10	F-Ne	0.30	21–25	Sc-Mn	0.05
11–12	Na-Mg	0.22	26–28	Fe-Ni	0.12

The spectrum of electrons and positrons incident at the top of the atmosphere is steeper than the spectra of protons and nuclei, as shown in Fig. 20.2. The positron fraction is about 10% in the region in which it is measured (< 20 GeV), but it is not yet fully understood [8].

Above 10 GeV the fraction of antiprotons to protons is about 10^{-4} , and there is evidence for the kinematic suppression at lower energy expected for secondary antiprotons [9,10,11,12]. There is at this time no evidence for a significant primary component of antiprotons.

20.2. Cosmic rays in the atmosphere

Figure 20.3 shows the vertical fluxes of the major cosmic ray components in the atmosphere in the energy region where the particles are most numerous (except for electrons, which are most numerous near their critical energy, which is about 81 MeV in air). Except for protons and electrons near the top of the atmosphere, all particles are produced in interactions of the primary cosmic rays in the air. Muons and neutrinos are products of the decay of charged mesons, while electrons and photons originate in decays of neutral mesons.




Figure 20.1: Major components of the primary cosmic radiation (from Ref. 1).

Most measurements are made at ground level or near the top of the atmosphere, but there are also measurements of muons and electrons from airplanes and balloons. Fig. 20.3 includes recent measurements of negative muons [3,13,14,15]. Since $\mu^+(\mu^-)$ are produced in association with $\nu_\mu(\bar{\nu}_\mu)$, the measurement of muons near the maximum of the intensity curve for the parent pions serves to calibrate the atmospheric ν_μ beam [16].

4 20. Cosmic rays




Figure 20.2: Differential spectrum of electrons plus positrons multiplied by E^3 (data summary from Ref. 7). The dashed line shows the proton spectrum multiplied by 0.01.




Figure 20.3: Vertical fluxes of cosmic rays in the atmosphere with $E > 1$ GeV estimated from the nucleon flux of Eq. (20.2). The points show measurements of negative muons with $E_\mu > 1$ GeV [3,13,14,15].

Because muons typically lose almost two GeV in passing through the atmosphere, the

comparison near the production altitude is important for the sub-GeV range of $\nu_\mu(\bar{\nu}_\mu)$ energies.

The flux of cosmic rays through the atmosphere is described by a set of coupled cascade equations with boundary conditions at the top of the atmosphere to match the primary spectrum. Numerical or Monte Carlo calculations are needed to account accurately for decay and energy-loss processes, and for the energy-dependences of the cross sections and of the primary spectral index γ . Approximate analytic solutions are, however, useful in limited regions of energy [17]. For example, the vertical intensity of nucleons at depth X (g cm^{-2}) in the atmosphere is given by

$$I_N(E, X) \approx I_N(E, 0) e^{-X/\Lambda}, \quad (20.3)$$

where Λ is the attenuation length of nucleons in air.

The corresponding expression for the vertical intensity of charged pions with energy $E_\pi \ll \epsilon_\pi = 115$ GeV is

$$I_\pi(E_\pi, X) \approx \frac{Z_{N\pi}}{\lambda_N} I_N(E_\pi, 0) e^{-X/\Lambda} \frac{X E_\pi}{\epsilon_\pi}. \quad (20.4)$$

This expression has a maximum at $t = \Lambda \approx 120$ g cm^{-2} , which corresponds to an altitude of 15 kilometers. The quantity $Z_{N\pi}$ is the spectrum-weighted moment of the inclusive distribution of charged pions in interactions of nucleons with nuclei of the atmosphere. The intensity of low-energy pions is much less than that of nucleons because $Z_{N\pi} \approx 0.079$ is small and because most pions with energy much less than the critical energy ϵ_π decay rather than interact.

20.3. Cosmic rays at the surface

20.3.1. Muons: Muons are the most numerous charged particles at sea level (see Fig. 20.3). Most muons are produced high in the atmosphere (typically 15 km) and lose about 2 GeV to ionization before reaching the ground. Their energy and angular distribution reflect a convolution of production spectrum, energy loss in the atmosphere, and decay. For example, 2.4 GeV muons have a decay length of 15 km, which is reduced to 8.7 km by energy loss. The mean energy of muons at the ground is ≈ 4 GeV. The energy spectrum is almost flat below 1 GeV, steepens gradually to reflect the primary spectrum in the 10–100 GeV range, and steepens further at higher energies because pions with $E_\pi > \epsilon_\pi \approx 115$ GeV tend to interact in the atmosphere before they decay. Asymptotically ($E_\mu \gg 1$ TeV), the energy spectrum of atmospheric muons is one power steeper than the primary spectrum. The integral intensity of vertical muons above 1 GeV/ c at sea level is ≈ 70 $\text{m}^{-2}\text{s}^{-1}\text{sr}^{-1}$ [18,19]. Experimentalists are familiar with this number in the form $I \approx 1$ $\text{cm}^{-2} \text{min}^{-1}$ for horizontal detectors.

The overall angular distribution of muons at the ground is $\propto \cos^2 \theta$, which is characteristic of muons with $E_\mu \sim 3$ GeV. At lower energy the angular distribution becomes increasingly steeper, while at higher energy it flattens and approaches a $\sec \theta$ distribution for $E_\mu \gg \epsilon_\pi$ and $\theta < 70^\circ$.

6 20. Cosmic rays




Figure 20.4: Spectrum of muons at $\theta = 0^\circ$ (\blacklozenge [18], \blacksquare [20], \blacktriangledown [21], \blacktriangle [22], X,+ [23]), and $\theta = 75^\circ$ \diamond [24]).

Figure 20.4 shows the muon energy spectrum at sea level for two angles. At large angles low energy muons decay before reaching the surface and high energy pions decay before they interact, thus the average muon energy increases. An approximate extrapolation formula valid when muon decay is negligible ($E_\mu > 100/\cos\theta$ GeV) and the curvature of the Earth can be neglected ($\theta < 70^\circ$) is

$$\frac{dN_\mu}{dE_\mu} \approx \frac{0.14 E_\mu^{-2.7}}{\text{cm}^2 \text{ s sr GeV}} \times \left\{ \frac{1}{1 + \frac{1.1 E_\mu \cos\theta}{115 \text{ GeV}}} + \frac{0.054}{1 + \frac{1.1 E_\mu \cos\theta}{850 \text{ GeV}}} \right\}, \quad (20.5)$$

where the two terms give the contribution of pions and charged kaons. Eq. (20.5) neglects a small contribution from charm and heavier flavors which is negligible except at very high energy [27].

The muon charge ratio reflects the excess of π^+ over π^- in the forward fragmentation region of proton initiated interactions together with the fact that there are more protons than neutrons in the primary spectrum. The charge ratio is between 1.1 and 1.4 from 1 GeV to 100 GeV [18,23]. Below 1 GeV there is a systematic dependence on location due to geomagnetic effects. [23]

20.3.2. Electromagnetic component: At the ground, this component consists of electrons, positrons, and photons primarily from electromagnetic cascades initiated by decay of neutral and charged mesons. Muon decay is the dominant source of low-energy electrons at sea level. Decay of neutral pions is more important at high altitude or when the energy threshold is high. Knock-on electrons also make a small contribution at low energy [25]. The integral vertical intensity of electrons plus positrons is very approximately 30, 6, and $0.2 \text{ m}^{-2}\text{s}^{-1}\text{sr}^{-1}$ above 10, 100, and 1000 MeV respectively [19,26], but the exact numbers depend sensitively on altitude, and the angular dependence is complex because of the different altitude dependence of the different sources of electrons [25,26,28]. The ratio of photons to electrons plus positrons is approximately 1.3 above a GeV and 1.7 below the critical energy [28].

20.3.3. Protons: Nucleons above $1 \text{ GeV}/c$ at ground level are degraded remnants of the primary cosmic radiation. The intensity is approximately represented by Eq. (20.3) with the replacement $t \rightarrow t/\cos\theta$ for $\theta < 70^\circ$ and an attenuation length $\Lambda = 123 \text{ g cm}^{-2}$. At sea level, about 1/3 of the nucleons in the vertical direction are neutrons (up from $\approx 10\%$ at the top of the atmosphere as the n/p ratio approaches equilibrium). The integral intensity of vertical protons above $1 \text{ GeV}/c$ at sea level is $\approx 0.9 \text{ m}^{-2}\text{s}^{-1}\text{sr}^{-1}$ [19,29].

20.4. Cosmic rays underground

Only muons and neutrinos penetrate to significant depths underground. The muons produce tertiary fluxes of photons, electrons, and hadrons.

20.4.1. Muons: As discussed in Section 23.6 of this *Review*, muons lose energy by ionization and by radiative processes: bremsstrahlung, direct production of e^+e^- pairs, and photonuclear interactions. The total muon energy loss may be expressed as a function of the amount of matter traversed as

$$-\frac{dE_\mu}{dX} = a + bE_\mu, \quad (20.6)$$

where a is the ionization loss and b is the fractional energy loss by the three radiation processes. Both are slowly varying functions of energy. The quantity $\epsilon \equiv a/b$ ($\approx 500 \text{ GeV}$ in standard rock) defines a critical energy below which continuous ionization loss is more important than radiative losses. Table 20.2 shows a and b values for standard rock as a function of muon energy. The second column of Table 20.2 shows the muon range in standard rock ($A = 22$, $Z = 11$, $\rho = 2.65 \text{ g cm}^{-3}$). These parameters are quite sensitive to the chemical composition of the rock, which must be evaluated for each experimental location.

The intensity of muons underground can be estimated from the muon intensity in the atmosphere and their rate of energy loss. To the extent that the mild energy dependence of a and b can be neglected, Eq. (20.6) can be integrated to provide the following relation between the energy $E_{\mu,0}$ of a muon at production in the atmosphere and its average energy E_μ after traversing a thickness X of rock (or ice or water):

$$E_\mu = (E_{\mu,0} + \epsilon)e^{-bX} - \epsilon. \quad (20.7)$$

8 20. Cosmic rays

Table 20.2: Average muon range R and energy loss parameters calculated for standard rock[30]. Range is given in km-water-equivalent, or 10^5 g cm^{-2} .

E_μ GeV	R km.w.e.	a MeV g $^{-1}$ cm 2	b_{brems} ————	b_{pair} $10^{-6} \text{ g}^{-1} \text{ cm}^2$	b_{nucl} $10^{-6} \text{ g}^{-1} \text{ cm}^2$	$\sum b_i$ ————
10	0.05	2.17	0.70	0.70	0.50	1.90
100	0.41	2.44	1.10	1.53	0.41	3.04
1000	2.45	2.68	1.44	2.07	0.41	3.92
10000	6.09	2.93	1.62	2.27	0.46	4.35

Especially at high energy, however, fluctuations are important and an accurate calculation requires a simulation that accounts for stochastic energy-loss processes [31].

There are two depth regimes for Eq. (20.7). For $X \ll b^{-1} \approx 2.5$ km water equivalent, $E_{\mu,0} \approx E_\mu(X) + aX$, while for $X \gg b^{-1}$ $E_{\mu,0} \approx (\epsilon + E_\mu(X)) \exp(bX)$. Thus at shallow depths the differential muon energy spectrum is approximately constant for $E_\mu < aX$ and steepens to reflect the surface muon spectrum for $E_\mu > aX$, whereas for $X > 2.5$ km.w.e. the differential spectrum underground is again constant for small muon energies but steepens to reflect the surface muon spectrum for $E_\mu > \epsilon \approx 0.5$ TeV. In the deep regime the shape is independent of depth although the intensity decreases exponentially with depth. In general the muon spectrum at slant depth X is

$$\frac{dN_\mu(X)}{dE_\mu} = \frac{dN_\mu}{dE_{\mu,0}} \frac{dE_{\mu,0}}{dE_\mu} = \frac{dN_\mu}{dE_{\mu,0}} e^{bX}, \quad (20.8)$$

where $E_{\mu,0}$ is the solution of Eq. (20.7) in the approximation neglecting fluctuations.

Fig. 20.5 shows the vertical muon intensity versus depth. In constructing this “depth-intensity curve,” each group has taken account of the angular distribution of the muons in the atmosphere, the map of the overburden at each detector, and the properties of the local medium in connecting measurements at various slant depths and zenith angles to the vertical intensity. Use of data from a range of angles allows a fixed detector to cover a wide range of depths. The flat portion of the curve is due to muons produced locally by charged-current interactions of ν_μ .

20.4.2. Neutrinos: Because neutrinos have small interaction cross sections, measurements of atmospheric neutrinos require a deep detector to avoid backgrounds. There are two types of measurements: contained (or semi-contained) events, in which the vertex is determined to originate inside the detector, and neutrino-induced muons. The latter are muons that enter the detector from zenith angles so large (*e.g.*, nearly horizontal or upward) that they cannot be muons produced in the atmosphere. In neither case is the neutrino flux measured directly. What is measured is a convolution of the neutrino flux and cross section with the properties of the detector (which includes the surrounding medium in the case of entering muons).




Figure 20.5: Vertical muon intensity vs depth (1 km.w.e. = 10^5 g cm $^{-2}$ of standard rock). The experimental data are from: \diamond : the compilations of Crouch [32], \square : Baksan [33], \circ : LVD [34], \bullet : MACRO [35], \blacksquare : Frejus [36]. The shaded area at large depths represents neutrino induced muons of energy above 2 GeV. The upper line is for horizontal neutrino-induced muons, the lower one for vertically upward muons.

Contained and semi-contained events reflect neutrinos in the sub-GeV to multi-GeV region where the product of increasing cross section and decreasing flux is maximum. In the GeV region the neutrino flux and its angular distribution depend on the geomagnetic location of the detector and, to a lesser extent, on the phase of the solar cycle. Naively, we expect $\nu_\mu/\nu_e = 2$ from counting neutrinos of the two flavors coming from the chain of pion and muon decay. This ratio is only slightly modified by the details of the decay kinematics, but the fraction of electron neutrinos gradually decreases above a GeV as parent muons begin to reach the ground before decaying. Experimental measurements have to account for the ratio of $\bar{\nu}/\nu$, which have cross sections different by a factor of 3 in this energy range. In addition, detectors generally have different efficiencies for detecting muon neutrinos and electron neutrinos which need to be accounted for in comparing

10 20. Cosmic rays

measurements with expectation. Fig. 20.6 shows the distributions of the visible energy in the Super-Kamiokande detector [39] for electron-like and muon-like charged current neutrino interactions. Contrary to expectation, the numbers of the two classes of events are similar rather than different by a factor of two. The exposure for the data sample shown here is 50 kiloton-years. The falloff of the muon-like events at high energy is a consequence of the poor containment for high energy muons. Corrections for detection efficiencies and backgrounds are, however, insufficient to account for the large difference from the expectation.




Figure 20.6: Sub-GeV and multi-GeV neutrino interactions from SuperKamiokande [39]. The plot shows the spectra of visible energy in the detector.

Two well-understood properties of atmospheric cosmic rays provide a standard for comparison of the measurements of atmospheric neutrinos. These are the “sec θ effect” and the “east-west effect”. The former refers originally to the enhancement of the flux of > 10 GeV muons (and neutrinos) at large zenith angles because the parent pions propagate more in the low density upper atmosphere where decay is enhanced relative to interaction. For neutrinos from muon decay, the enhancement near the horizontal becomes important for $E_\nu > 1$ GeV and arises mainly from the increased pathlength through the atmosphere for muon decay in flight. Fig. 20.7 from Ref. 40 shows a comparison between measurement and expectation for the zenith angle dependence of multi-GeV electron-like (mostly ν_e) and muon-like (mostly ν_μ) events separately. The ν_e show an enhancement near the horizontal and approximate equality for nearly upward ($\cos \theta \approx -1$) and nearly downward ($\cos \theta \approx 1$) events. There is, however, a very significant deficit of upward ($\cos \theta < 0$) ν_μ events, which have long pathlengths comparable to the radius of the Earth. This pattern has been interpreted as evidence for oscillations involving muon

neutrinos [39]. (See the article on neutrino properties in this *Review*.) Including three dimensional effects in the calculation of atmospheric neutrinos may change somewhat the expected angular distributions of neutrinos at low energy [41], but it does not change the fundamental expectation of up-down symmetry, which is the basis of the evidence for oscillations.




Figure 20.7: Zenith-angle dependence of multi-GeV neutrino interactions from SuperKamiokande [40]. The shaded boxes show the expectation in the absence of any oscillations. The lines show fits with some assumed oscillation parameters, as described in Ref. 40.

The east-west effect [42,43] is the enhancement, especially at low geomagnetic latitudes, of cosmic rays incident on the atmosphere from the west as compared to those from the east. This is a consequence of the fact that the cosmic rays are positively charged nuclei which are bent systematically in one sense in the geomagnetic field. Not all trajectories can reach the atmosphere from outside the geomagnetic field. The standard procedure to see which trajectories are allowed is to inject antiprotons outward from near the top of the atmosphere in various directions and see if they escape from the geomagnetic field without becoming trapped indefinitely or intersecting the surface of the Earth. Any direction in which an antiproton of a given momentum can escape is an allowed direction from which a proton of the opposite momentum can arrive. Since the geomagnetic field is oriented from south to north in the equatorial region, antiprotons injected toward the east are bent back towards the Earth. Thus there is a range of momenta and zenith angles for which positive particles cannot arrive from the east but can arrive from the west. This east-west asymmetry of the incident cosmic rays induces a similar asymmetry on the secondaries, including neutrinos. Since this is an azimuthal effect, the resulting asymmetry is independent of possible oscillations, which depend on pathlength (equivalently zenith angle), but not on azimuth. Fig. 20.8 (from Ref. 44) is a comparison of data and expectation for this effect, which serves as a consistency check of the measurement and analysis.

Muons that enter the detector from outside after production in charged-current interactions of neutrinos naturally reflect a higher energy portion of the neutrino

12 20. Cosmic rays

Table 20.3: Measured fluxes ($10^{-13} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$) of neutrino-induced muons as a function of the effective minimum muon energy E_μ .

$E_\mu >$	1 GeV	1 GeV	1 GeV	2 GeV	3 GeV	3 GeV
Ref.	CWI [45]	Baksan [46]	MACRO [47]	IMB [48]	Kam [49]	SuperK [50]
F_μ	2.17 ± 0.21	2.77 ± 0.17	2.29 ± 0.15	2.26 ± 0.11	1.94 ± 0.12	1.74 ± 0.07

spectrum than contained events because the muon range increases with energy as well as the cross section. The relevant energy range is $\sim 10 < E_\nu < 1000 \text{ GeV}$, depending somewhat on angle. Neutrinos in this energy range show a $\sec \theta$ effect similar to muons (see Eq. (20.5)). This causes the flux of horizontal neutrino induced muons to be approximately a factor two higher than the vertically upward flux. The upper and lower edges of the horizontal shaded region in Fig. 20.5 correspond to horizontal and vertical intensities of neutrino-induced muons. Table 20.3 gives the measured fluxes of upward-moving neutrino-induced muons averaged over the lower hemisphere. Generally the definition of minimum muon energy depends on where it passes through the detector. The tabulated effective minimum energy estimates the average over various accepted trajectories.

20.5. Air showers

So far we have discussed inclusive or uncorrelated fluxes of various components of the cosmic radiation. An air shower is caused by a single cosmic ray with energy high enough for its cascade to be detectable at the ground. The shower has a hadronic core, which acts as a collimated source of electromagnetic subshowers, generated mostly from $\pi^0 \rightarrow \gamma\gamma$. The resulting electrons and positrons are the most numerous particles in the shower. The number of muons, produced by decays of charged mesons, is an order of magnitude lower.

Air showers spread over a large area on the ground, and arrays of detectors operated for long times are useful for studying cosmic rays with primary energy $E_0 > 100 \text{ TeV}$, where the low flux makes measurements with small detectors in balloons and satellites difficult.

Greisen [51] gives the following approximate expressions for the numbers and lateral distributions of particles in showers at ground level. The total number of muons N_μ with energies above 1 GeV is

$$N_\mu(> 1 \text{ GeV}) \approx 0.95 \times 10^5 \left(N_e / 10^6 \right)^{3/4}, \quad (20.9)$$

where N_e is the total number of charged particles in the shower (not just e^\pm). The number of muons per square meter, ρ_μ , as a function of the lateral distance r (in meters) from the center of the shower is

$$\rho_\mu = \frac{1.25 N_\mu}{2\pi \Gamma(1.25)} \left(\frac{1}{320} \right)^{1.25} r^{-0.75} \left(1 + \frac{r}{320} \right)^{-2.5}, \quad (20.10)$$




Figure 20.8: Azimuthal dependence of \sim GeV neutrino interactions from SuperKamiokande [44]. The cardinal points of the compass are S, E, N, W starting at 0. These are the direction from which the particles arrive. The lines show the expectation based on two different calculations, as described in Ref. 44.

where Γ is the gamma function. The number density of charged particles is

$$\rho_e = C_1(s, d, C_2) x^{(s-2)} (1+x)^{(s-4.5)} (1+C_2 x^d) . \quad (20.11)$$

Here s , d , and C_2 are parameters in terms of which the overall normalization constant $C_1(s, d, C_2)$ is given by

$$C_1(s, d, C_2) = \frac{N_e}{2\pi r_1^2} [B(s, 4.5 - 2s) + C_2 B(s + d, 4.5 - d - 2s)]^{-1} , \quad (20.12)$$

where $B(m, n)$ is the beta function. The values of the parameters depend on shower size (N_e), depth in the atmosphere, identity of the primary nucleus, etc. For showers with $N_e \approx 10^6$ at sea level, Greisen uses $s = 1.25$, $d = 1$, and $C_2 = 0.088$. Finally, x is r/r_1 , where r_1 is the Molière radius, which depends on the density of the atmosphere and hence on the altitude at which showers are detected. At sea level $r_1 \approx 78$ m. It increases with altitude.

The lateral spread of a shower is determined largely by Coulomb scattering of the many low-energy electrons and is characterized by the Molière radius. The lateral spread




Figure 20.9: The all-particle spectrum: \square [52], \blacktriangle [54], \blacktriangledown [55], ∇ [56], \triangle [57], $+$ [58], \times [59], \blacklozenge [60]. References for the high energy portion of the spectrum are given in Fig. 20.10.

of the muons (ρ_μ) is larger and depends on the transverse momenta of the muons at production as well as multiple scattering.

There are large fluctuations in development from shower to shower, even for showers of the same energy and primary mass—especially for small showers, which are usually well past maximum development when observed at the ground. Thus the shower size N_e and primary energy E_0 are only related in an average sense, and even this relation depends on depth in the atmosphere. One estimate of the relation is [52]

$$E_0 \sim 3.9 \times 10^6 \text{ GeV} (N_e/10^6)^{0.9} \quad (20.13)$$

for vertical showers with $10^{14} < E < 10^{17}$ eV at 920 g cm^{-2} (965 m above sea level). Because of fluctuations, N_e as a function of E_0 is not the inverse of Eq. (20.13). As E_0 increases the shower maximum (on average) moves down into the atmosphere and the relation between N_e and E_0 changes. At the maximum of shower development, there are approximately 2/3 particles per GeV of primary energy.

Detailed simulations and cross-calibrations between different types of detectors are necessary to establish the primary energy spectrum from air-shower experiments [52,53]. Figure 20.9 shows the “all-particle” spectrum. In establishing this spectrum, efforts have been made to minimize the dependence of the analysis on the primary composition. In the energy range above 10^{17} eV, the Fly’s Eye technique [71] is particularly useful because it can establish the primary energy in a model-independent way by observing most of the longitudinal development of each shower, from which E_0 is obtained by integrating the energy deposition in the atmosphere.

In Fig. 20.9 the differential energy spectrum has been multiplied by $E^{2.7}$ in order to display the features of the steep spectrum that are otherwise difficult to discern. The steepening that occurs between 10^{15} and 10^{16} eV is known as the *knee* of the spectrum. The feature between 10^{18} and 10^{19} eV is called the *ankle* of the spectrum.

If the cosmic ray spectrum below 10^{18} eV is of galactic origin, the *knee* could reflect the fact that some (but not all) cosmic accelerators have reached their maximum energy. Some types of expanding supernova remnants, for example, are estimated not to be able to accelerate particles above energies in the range of 10^{15} eV total energy per particle. Effects of propagation and confinement in the galaxy [61] also need to be considered.

The *ankle* has the classical characteristic shape [62] of a higher energy population of particles overtaking a lower energy population. A possible interpretation is that the higher energy population represents cosmic rays of extragalactic origin. If this is the case and if the cosmic rays are cosmological in origin, then there should be a cutoff around 5×10^{19} eV, resulting from interactions with the microwave background [63,64]. It is therefore of special interest that several events have been assigned energies above 10^{20} eV [65,66,67,68].

Figure 20.10 gives an expanded view of the high energy end of the spectrum. Not included in the figure are the preliminary results of the Hi-Res Fly's Eye group [69]. Their spectrum is consistent with the others and includes several events above 10^{20} eV. To expand the scale in Fig. 20.10, the differential flux is multiplied by E^3 , a process which amplifies small systematic differences in energy assignments into sizeable differences in rate. The different measurements agree with each other within 20% systematic shifts in energy scale and consistently show that the spectrum continues well past the expected cutoff for a cosmological distribution of sources. The implication is that some sources of the highest energy particles must be relatively nearby. For example, the attenuation length for protons at 2×10^{20} eV is 30 Mpc [70].




Figure 20.10: Expanded view of the highest energy portion of the cosmic-ray spectrum: \circ [71] (stereo), \diamond [71] (monocular) \blacksquare [72], \blacktriangledown [73].

References:

1. J.A. Simpson, *Ann. Rev. Nucl. and Part. Sci.* **33**, 323 (1983).
2. M. Boezio *et al.*, *Astrophys. J.* **518**, 457 (1999).
3. R. Bellotti *et al.*, *Phys. Rev.* **D60**, 052002 (1999).
4. AMS Collaboration, *Phys. Lett.* **B** (to be published, 1999).
5. T. Sanuki *et al.*, in *Proc. 26th Int. Cosmic Ray Conf.*, Salt Lake City, **3**, 93 (1999).
6. J.J. Engelmann *et al.*, *Astron. & Astrophys.* **233**, 96 (1990);
See also *Cosmic Abundances of Matter* (ed. C. Jake Waddington) A.I.P. Conf. Proceedings No. 183 (1988) p. 111.
7. S.W. Barwick *et al.*, *Astrophys. J.* **498**, 779 (1998).
8. Stephane Coutu *et al.*, *Astropart. Phys.* **11**, 429 (1999).
9. J.W. Mitchell *et al.*, *Phys. Rev. Lett.* **76**, 3057 (1996).
10. M. Boezio *et al.*, *Astrophys. J.* **487**, 415 (1997).
11. M. Hof *et al.*, *Astrophys. J.* **467**, L33 (1996) See also G. Basini *et al.*, *Proc. 26th Int. Cosmic Ray Conf.*, Salt Lake City, **3**, 101 (1999).
12. H. Matsunaga *et al.*, *Phys. Rev. Lett.* **81**, 4052 (1998). See also S. Orito *et al.*, [astro-ph/9906426](https://arxiv.org/abs/astro-ph/9906426).
13. R. Bellotti *et al.*, *Phys. Rev.* **D53**, 35 (1996).
14. M. Boezio *et al.*, *Phys. Rev. Lett.* **82**, 4757 (1999).
15. S. Coutu *et al.*, *Proc. 26th Int. Cosmic Ray Conf.*, Salt Lake City, **2**, 68 (1999).
16. D.H. Perkins, *Astropart. Phys.* **2**, 249 (1994).
17. T.K. Gaisser, *Cosmic Rays and Particle Physics*, Cambridge University Press (1990).
18. M.P. De Pascale *et al.*, *J. Geophys. Res.* **98**, 3501 (1993).
19. K. Allkofer and P.K.F. Grieder, *Cosmic Rays on Earth*, Fachinformationszentrum, Karlsruhe (1984).
20. O.C. Allkofer, K. Carstensen, and W.D. Dau, *Phys. Lett.* **B36**, 425 (1971).
21. B.C. Rastin, *J. Phys.* **G10**, 1609 (1984).
22. C.A. Ayre *et al.*, *J. Phys.* **G1**, 584 (1975).
23. J. Kremer *et al.*, *Phys. Rev. Lett.* **83**, 4241 (1999).
24. H. Jokisch *et al.*, *Phys. Rev.* **D19**, 1368 (1979).
25. S. Hayakawa, *Cosmic Ray Physics*, Wiley, Interscience, New York (1969).
26. R.R. Daniel and S.A. Stephens, *Revs. Geophysics & Space Sci.* **12**, 233 (1974).
27. F. Halzen, R. Vázquez, and E. Zas, *Astropart. Phys.* **1**, 297 (1993).
28. K.P. Beuermann and G. Wibberenz, *Can. J. Phys.* **46**, S1034 (1968).
29. I.S. Diggory *et al.*, *J. Phys.* **A7**, 741 (1974).
30. D.E. Groom, N.V. Mokhov, and S.I. Striganov, "Muon stopping-power and range tables," *Atomic Data and Nuclear Data Tables*, to be published (2000).

18 20. Cosmic rays

31. P. Lipari and T. Stanev, *Phys. Rev.* **D44**, 3543 (1991).
32. M. Crouch, in *Proc. 20th Int. Cosmic Ray Conf.*, Moscow, **6**, 165 (1987).
33. Yu.M. Andreev, V.I. Gurentzov, and I.M. Kogai, in *Proc. 20th Int. Cosmic Ray Conf.*, Moscow, **6**, 200 (1987).
34. M. Aglietta *et al.*, (LVD Collaboration), *Astropart. Phys.* **3**, 311 (1995).
35. M. Ambrosio *et al.*, (MACRO Collaboration), *Phys. Rev.* **D52**, 3793 (1995).
36. Ch. Berger *et al.*, (Frejus Collaboration), *Phys. Rev.* **D40**, 2163 (1989).
37. K.S. Hirata *et al.*, (Kam-II Collaboration), *Phys. Lett.* **B280**, 146 (1992);
Y. Fukuda *et al.*, *Phys. Lett.* **B335**, 237 (1994).
38. R. Becker-Szendy *et al.*, (IMB Collaboration), *Phys. Rev.* **D46**, 3720 (1992);
See also D. Casper *et al.*, *Phys. Rev. Lett.* **66**, 2561 (1991).
39. Y. Fukuda *et al.*, *Phys. Rev. Lett.* **81**, 1562 (1998).
40. K. Scholberg *et al.*, in *Proc. Eighth Int. Workshop on Neutrino Telescopes* (ed. Milla Baldo Ceolin, edizioni papergraf, Venezia, 1999) p. 183.
41. G. Battistoni *et al.*, *Astropart. Phys.* **12**, 315 (2000).
42. T.H. Johnson, *Phys. Rev.* **43**, 834 (1933). See also T.H. Johnson & E.C. Street, *Phys. Rev.* **44**, 125 (1933).
43. L.W. Alvarez and A.H. Compton, *Phys. Rev.* **43**, 835 (1933).
44. T. Futagami *et al.*, *Phys. Rev. Lett.* **82**, 5194 (1999).
45. F. Reines *et al.*, *Phys. Rev. Lett.* **15**, 429 (1965).
46. M.M. Boliev *et al.*, in *Proceedings 3rd Int. Workshop on Neutrino Telescopes* (ed. Milla Baldo Ceolin), 235 (1991).
47. M. Ambrosio *et al.*, (MACRO) *Phys. Lett.* **B434**, 451 (1998). The number quoted for MACRO is the average over 90% of the lower hemisphere, $\cos\theta < -0.1$; see F. Ronga *et al.*, hep-ex/9905025.
48. R. Becker-Szendy *et al.*, *Phys. Rev. Lett.* **69**, 1010 (1992);
Proc. 25th Int. Conf. High-Energy Physics, Singapore (ed. K.K. Phua and Y. Yamaguchi, World Scientific, 1991) p. 662.
49. S. Hatakeyama *et al.*, *Phys. Rev. Lett.* **81**, 2016 (1998).
50. Y. Fukuda *et al.*, *Phys. Rev. Lett.* **82**, 2644 (1999).
51. K. Greisen, *Ann. Rev. Nucl. Sci.* **10**, 63 (1960).
52. M. Nagano *et al.*, *J. Phys.* **G10**, 1295 (1984).
53. M. Teshima *et al.*, *J. Phys.* **G12**, 1097 (1986).
54. N.L. Grigorov *et al.*, *Yad. Fiz.* **11**, 1058 (1970) and *Proc. 12th Int. Cosmic Ray Conf.*, Hobart, **2**, 206 (1971).
55. K. Asakimori *et al.*, *Proc. 23rd Int. Cosmic Ray Conf.*, Calgary, **2**, 25 (1993);
Proc. 22nd Int. Cosmic Ray Conf., Dublin, **2**, 57 and 97 (1991).

56. T.V. Danilova *et al.*, *Proc. 15th Int. Cosmic Ray Conf.*, Plovdiv, **8**, 129 (1977).
57. Yu. A. Fomin *et al.*, *Proc. 22nd Int. Cosmic Ray Conf.*, Dublin, **2**, 85 (1991).
58. M. Amenomori *et al.*, *Astrophys. J.* **461**, 408 (1996).
59. A. Roehring *et al.*, *Proc. 26th Int. Cosmic Ray Conf.*, Salt Lake City, **1**, 214 (1999).
60. M.A.K. Glasmacher *et al.*, *Astropart. Phys.* **10**, 291 (1999).
61. V.S. Ptuskin *et al.*, *Astron. & Astrophys.* **268**, 726 (1993).
62. B. Peters, *Nuovo Cimento* **22**, 800 (1961).
63. K. Greisen, *Phys. Rev. Lett.* **16**, 748 (1966).
64. G.T. Zatsepin and V.A. Kuz'min, *Sov. Phys. JETP Lett.* **4**, 78 (1966).
65. J. Linsley, *Phys. Rev. Lett.* **10**, 146 (1963).
66. D.J. Bird *et al.*, *Astrophys. J.* **441**, 144 (1995).
67. N. Hayashima *et al.*, *Phys. Rev. Lett.* **73**, 3941 (1994).
68. A.A. Watson, *Proc. of the 1994 Summer Study on Nuclear and Particle Astrophysics and Cosmology for the Next Millenium*, Snowmass CO, 1994, ed. by E.W. Kolb *et al.* (World Scientific, Singapore, to be published, 1996).
69. C. Jui *et al.*, highlight talk, *26th International Cosmic Ray Conf.*, Salt Lake City, 1999, to be published in A.I.P. Conference Proceedings.
70. V.S. Berezhinskii *et al.*, *Astrophysics of Cosmic Rays*, North-Holland (1990).
71. D.J. Bird *et al.*, *Astrophys. J.* **424**, 491 (1994).
72. M. Takeda *et al.*, *Phys. Rev. Lett.* **81**, 1163 (1998).
73. M.A. Lawrence, R.J.O. Reid, and A.A. Watson, *J. Phys.* **G17**, 733 (1991).